



Zinc powder 15%.

Product usable in Organic Agriculture according to Regulation (EU) No. 2018/848 and 2021/1165.  
Product that complies with NOP (National Organic Program - USDA)

### Description- specifications

**FERTICELL ZINC 15%** is an ecological fertilizer, used for both foliar applications and fertigation. It is used to prevent and correct zinc deficiencies, being a cofactor of many key enzyme processes, that favours protein synthesis as well as auxins synthesis by plants.

**FERTICELL ZINC 15%** is especially indicated to improve the flowering processes of crops, and to cover zinc deficiencies, which is one of the most common. Among the crops most sensitive to zinc deficiency, are corn, cotton, citrus, apple, peach, and avocado.

**FERTICELL ZINC 15%** is used in the vegetative phase of the crop, will act as a cell growth regulator.

### Uses



Foliar  
Fertigation

### Available containers



20 Kg

### Certifications



Product suitable for use in Organic Agriculture in accordance with Regulations (EU) n°2018/848 and 2021/1165, the NOP Regulation.

Control ECOCERT

F- 32600

Agroplasma SL  
Polígono Industrial Santa Teresa  
Calle Torre del Mar N°56  
29004 Málaga, Spain

### GUARANTEED ANALYSIS:

Composition	Percentage (w/w)	
Zinc (Zn), total	≥ 15 %	
Total Nitrogen (N)	≥ 8 %	

### PHYSICAL PROPERTIES

State: Powder

Colour: Grey

### GENERAL USAGE AND DOSAGE RECOMMENDATIONS

Crop	Foliar dose	Fertigation dose
<b>Fruit trees:</b> Citrus fruits, Apples, Pears, Blueberries, Almonds, Cherries, Peaches, Avocado, Mango, Banana, etc...	-Maintenance: 2-3 Kg/ha or 200-300 mg/100 l/ application -Strong deficiency: Up to 4 applications	
<b>Horticultural crops:</b> Tomato, Potato, Cucumber, Melon Courgette, Cauliflower, Onion, Carrot, Pepper, etc...	-Maintenance: 2-3 Kg/ha or 200-300 mg/100 l/ application -Strong deficiency: Up to 4 applications	
<b>Extensive crops:</b> Cereals, Alfalfa, Cotton, Beetroot, Corn, Sunflower, etc..	-Maintenance: 2-3 Kg/ha or 200-300 mg/100 l/ application -Strong deficiency: Up to 4 applications	

· The specified dosages are general recommendations. The amounts depend on the crop, phenological state, level of deficiency and soil type.

· The physical and appearance characteristics of the product may be altered due to the ecological nature of the product. Keep the product at room temperature, in a cool and dry place.

· Use under technical advice. Keep the product at **room temperature**.

#### Important notes:

1. Due to its high concentration in zinc, the product must be prediluted before being poured into the irrigation tank or spray.
2. In fertigation applications, divide input being as maximum: 3Kg/ ha, at intervals of 10-15 days.